## **NUCLEAR MEDICINE EXAM INFORMATION**

EXAM	EXAM PREPARATION	ESTIMATED EXAM DURATION
BONE & GALLIUM		
Bone Scan	No exam preparation	2 part exam (4 hours total)
Gallium Scan	No exam preparation	2 non-consecutive days
	Gallium injection (15 min) → return for scan (1+ hour) 2-3 days after in	njection
Osteomyelitis Scan	Bone scan will be done first and gallium scan will be booked as neede	ed after the bone scan
BRAIN & CEREBRAL SPINAL FLUID (CSF)		
Brain Perfusion Scan	No exam preparation	2 part exam (2 hours total)
Cavernous Sinus - RBC	No exam preparation	2 part exam (4 hours total)
VP Shunt Scan	No exam preparation	1½ hours
ENDOCRINE	· ·	
Iodine Scan – Diagnostic	No food or drink 2 hours before + 2 hours after the start of the test	2 non-consecutive days
-	lodine Wednesday (½ hour) → return on Friday for the scan (1+ hour)	_
	Follow your referring doctor's instructions for low iodine diet, thyrogen	
Iodine Scan – Post-Therapy	Follow your doctor's preparation instructions for iodine therapy	1 hour
Parathyroid Scan	No exam preparation	2 part exam (3 hours total)
Thyroid Uptake & Scan	No food or drink 2 hours before + 2 hours after the start of the test	2 back-to back days
- '	Ask your doctor about stopping thyroid medicines	Day 1: 2 parts (2½ hours total)
	No x-ray/CT IV contrast dye 6 weeks or cough medicines 4 weeks	Day 2: 1 hour
GASTRO-INTESTINAL	· · · · · · · · · · · · · · · · · · ·	•
C-14 Urea Breath Test	No food or drink 4 hours before the test	½ hour
	No antibiotics or Pepto-Bismol 1 month + no Sucralfate or proton pum	
Esophageal Transit	No food or drink 4 hours before the test	½ hour
Gastric Emptying Scan	No food or drink 4 hours before the test	4½ hours
, , J	Call us at 416-864-5115 if you have food allergies, sensitivities, or die	
	No motility, laxative, anti-diarrhea, or narcotic medicines 2 days + no	-
Gastro-Intestinal Bleed Scan	No food or drink 4 hours before the test	2-4 hours
Meckel's Diverticulum Scan	No food or drink 4 hours before the test	2-4 hours
Salivary Scan	No food, gum, candy, lozenges 2 hours before the test (only water)	1 hour
LIVER, SPLEEN, & GALLBLADDER		
Hepatobiliary (Biliary) Scan	No food or drink 4 hours before the test	2-4 hours
1 3 ( 3 /	No narcotic pain medicines 24 hours before the test	
Liver/Spleen – Colloid Scan	No exam preparation	1 hour
Liver/Spleen – RBC Scan	No exam preparation	2 part exam (4 hours total)
Splenic Sequestration Scan	No exam preparation	2 hours
LYMPHATIC (LYMPH NODE		
Sentinel Node – Breast	No exam preparation; follow surgical instructions as applicable	½ hour
Sentinel Node – Melanoma	No exam preparation; follow surgical instructions as applicable	1½ hours
RADIONUCLIDE THERAPY		
I-131 Radioiodine Therapy	No food or drink 2 hours before + 2 hours after the start of the test	½ hour
,	Follow your referring doctor's instructions for low iodine diet, thyrogen	
RESPIRATORY (LUNG)	, , , , , , , , , , , , , , , , , , , ,	, , , ,
Lung Scan (V/Q Scan)	No exam preparation; recent chest x-ray may be required	1 hour
Systemic Shunt Scan	No exam preparation	1 hour
Quantitative Lung Perfusion	No exam preparation	1 hour
URINARY (KIDNEYS)	* For all renal scans drink 4 glasses of water leading up to test t	
Renal Scan	Drink 4 glasses of water	1 hour
Renal Scan with Blood GFR	Drink 4 glasses of water; ask your doctor about stopping medicines	4 hours
Renal Scan with Captopril	Drink 4 glasses of water; no food 4 hours + no ACE inhibitors 48 hrs	2 hours
Renal Scan with Lasix	Drink 4 glasses of water; ask your doctor about stopping medicines	1½ hours
Renal Cortical Scan	Drink 4 glasses of water	2 part test (4 hours total)
		=

Exam durations are approximate and may vary. Preparation instructions are guidelines and may vary.
Call us at 416-864-5115 if you have questions about your test.

